



TES Experiment Overview & INTEX-B Validation Flights

10/27/2005

TES Validation Team

Jet Propulsion Laboratory, California Institute of Technology





TES Primary Science Goals



Measurements of Tropospheric O_3 and precursors that will significantly improve our understanding of:

- Intercontinental Transport of Pollutants
- Tropospheric Chemistry
- Global O₃ Sources & Sinks
- Role of O₃ in Climate Forcing/Feedback





INTEX-B



INTEX-B can provide:


- DC-8 aircraft measurements of tropospheric ozone, carbon monoxide and nitric acid.
- Significant number of profiles (remote and in-situ) to compare with TES data.
- → INTEX-B provides an ideal opportunity for obtaining data sets for validation and science analysis of TES data.
- → TES can provide global and regional data in areas of scientific interest during INTEX-B.





TES Global Survey (every other day): Separation of ~1.6° Between Nadir Scenes










TES Global Survey as indicator of O₃ transport



TES July 7 2005 (Upper Troposphere)



(image provided by John Worden)






TES Special Observations



- The following three types of TES special observations can be scheduled for non-global survey days:
- Step & Stare (O_3, H_2O, CO)
 - Series of 150 nadir observations spaced 5.1 seconds apart (~34 km along the Aura nadir track).
 - Covers approximate 60 degrees of latitude.
 - Can be started at any latitude.
- Limb (HNO₃)
 - Series of limb observations
 - Can be started at any latitude.
- Combined Step & Stare and **Limb** looking at the same air mass (under development)



Green: Global Survey Violet: Calibration Magenta: SO-Ascending Grange: SO-Descending





Validation Goals for INTEX-B



TES Validation Priorities:

- Ozone
 - Lidar profiles of ozone under different atmospheric conditions during long, level flight legs along Aura orbit track
 - In situ profiles of ozone along the Aura orbit track.
- Carbon Monoxide
 - Similar to ozone, getting in situ profiles under a variety of atmospheric conditions along the Aura flight track
- Nitric Acid (Measured in the Limb)
 - Long level legs between 9-11 km, preferably along the TES limb track.






Validation Lessons Learned During AVE & PAVE Campaigns



- Aircraft flights along the Aura track are preferable on days that TES can make special observations.
 - TES Global Survey days are extremely difficult to change.
- Aircraft profiles from maximum altitude all the way down to sea level are highly desirable for the DC-8 in situ measurements.
 - Meaningful comparisons of TES with in situ data are very difficult unless the entire tropospheric profiles are available.
- Ozonesondes:
 - These have been, and will continue to be, a very high priority for TES validation.
- Variability of water vapor makes comparisons between TES and aircraft difficult. The best approach for water validation may be a statistical comparison between TES, AIRS, and sondes.





TES Measurement Status



- Global Surveys (with limb scans) taken Sep 2004 Apr 2005
 - 73 16-orbit Global Surveys acquired with limb data
 - Nadir targets ~5° apart along orbit track (2 nadir scans averaged)
- Global Surveys without limb scans started May 2005
 - 3x more nadir targets, ~1.6° separation, no averaging of scenes
 - Limb mode still available for special observations, but removed from the Global Survey mode to preserve instrument lifetime.
 - No data for June 2005 since instrument was in safe mode.
- Step & Stare, Stare and Transect modes used for special observations
 - AVE (Oct 2004 Nov 2004)
 - PAVE (Jan 2005 Feb 2005)
 - S. Atlantic (Sep 2004 Oct 2004, Jan 2005 Feb 2005)
 - N. America (July 31, 2005 August 6, 2005)
 - Lake Tahoe: Stare mode for geolocation, radiance val. (Aug 05 Sep 05)
 - S. America: biomass burning transects (Aug 2005 Sep 2005)





TES Data Availability



- Current TES data at the Langley ASDC is a Beta version (v001)
 - Nadir scenes only.
 - L2 retrievals from special observations to start delivery Oct 2005.
 - Caveats listed for L1B calibration and L2 land retrieval issues.
 - Files in HDF format
- Data taken during AVE 04 and PAVE 05 (limited) is available at the Aura Validation Data Center (AVDC).
 - Data is in IDL "Save" file format
- Next release (v002) in Spring 2006
 - L1B calibration improved substantially & validated
 - Will include limb retrievals (with HNO3)
 - HDO product added for nadir views





Data Quality of TES Data (v001)



Level 1B:

- TES L1B data products have systematic errors that will be fixed in v002 data.
 - Systematic errors are estimated after radiometric calibration and show an average radiance error of approximately 2%.
 - Errors specific to each target spectrum are available within the data products.

Level 2:

- TES L2 products that are ready for scientific use are nadir retrievals of ozone, carbon monoxide, temperature and water for ocean target scenes.
- Caveats on TES L2 Products:
 - Land scenes are reported but should be used with caution due to a software bug.
 - TES L2 retrievals over the Sahara desert regions have problems with the a priori estimates of emissivity and should not be used.
 - Retrieved parameters poleward of 60 should be used with caution due to the low brightness temperatures associated with these scenes.
 - As a means of accounting for systematic errors in L1B radiances, current L2 profiles include the retrieval of a calibration scaling parameter.
- The error estimates included in the L2 data products are meaningful based on the current validation analysis.







Part of the research described here was carried out by the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA.

